

## Evaluation of antioxidant and antibacterial activities of extracts of *Celtis philippensis* Blanco (Ulmaceae) on multidrug resistant wound pathogens

Philip A. Idowu<sup>1</sup>, Faith J. Afolayan<sup>1</sup>, Charles O. Olaiya<sup>2</sup>

<sup>1</sup>Department of Pharmaceutical Microbiology, Faculty of Pharmacy, University of Ibadan, Nigeria.

<sup>2</sup>Department of Biochemistry, Faculty of Basic Medical Sciences, University of Ibadan, Nigeria

Corresponding author: Philip Adegboyega Idowu,  
E-mail: igboyega@yahoo.com; Phone: +2348033524399

### ABSTRACT

**Background:** Antimicrobial resistance among wound pathogens is increasing, creating the need to search for new and effective agents, especially among medicinal plants used to treat wounds traditionally. There is a worldwide search for medicinal plants with good antibacterial and antioxidant activities to treat infected wounds.

**Objectives:** *Celtis philippensis* used locally to treat infections was evaluated for its antioxidant and antibacterial activities against multidrug resistant (MDR) pathogens from infected wounds from three Nigerian hospitals.

**Methods:** Methanol extract of *C. philippensis* stem was tested at 20 and 10mg/ml on 32 wound pathogens comprising of *Staphylococcus aureus* (14), *Pseudomonas aeruginosa* (10) and *Klebsiella pneumoniae* (8). Agar diffusion methods was used to determine antibiogram and susceptibility to extracts while agar dilution method was used to determine minimum inhibitory concentrations (MIC) and minimum bactericidal concentrations (MBC). The antioxidant potential was evaluated by four methods: catalase activity, lipid peroxidation inhibition, hydrogen peroxide radical scavenging activity and DPPH radical scavenging activity. Statistical analysis was done using ANOVA.

**Results:** Antibiogram showed that the test organisms were MDR bacteria. The extract was active on all the tested pathogens with zones of inhibition of between 13-35mm. The MIC and MBC values ranged between 0.625-5mg/ml for all the test organisms and MIC index indicated bactericidal activity. The methanol extract showed good antioxidant activity (lipid peroxidation inhibition and catalase activity) comparable with ascorbic acid standard.

**Conclusion:** Antioxidant and antibacterial activities of *Celtis philippensis* justified its folkloric use in treating infections. Moreover, good activity on multidrug resistant wound pathogens highlights its potentials as a source of antimicrobial agents to treat wounds infected with MDR pathogens.

**Keywords:** *Celtis philippensis*, multidrug resistant pathogens, antibacterial, antioxidant, wounds.

## Évaluation des activités antioxydantes et antibactériennes d'extraits de *Celtis philippensis* Blanco (Ulmaceae) sur des pathogènes multirésistants aux plaies

Philip A. Idowu<sup>1</sup>, Faith J. Afolayan<sup>1</sup>, Charles O. Olaiya<sup>2</sup>

<sup>1</sup>Département de microbiologie pharmaceutique, Faculté de pharmacie, Université d'Ibadan, Nigéria.

<sup>2</sup>Département de la biochimie, Faculté des sciences médicales fondamentales, Université d'Ibadan, Nigéria

Auteur correspondant : Philip Adegboyega Idowu,  
E-mail : igboyega@yahoo.com; Téléphone : +2348033524399

### RESUME

**Contexte :** La résistance aux antimicrobiens parmi les agents pathogènes des plaies augmente, créant le besoin de rechercher de nouveaux agents efficaces, en particulier parmi les plantes médicinales utilisées traditionnellement pour traiter les plaies. Il existe une recherche mondiale de plantes médicinales ayant de bonnes activités antibactériennes et antioxydantes pour traiter les plaies infectées

**Objectifs :** Le *Celtis philippensis* utilisé localement pour traiter les infections a été évalué pour ses activités antioxydantes et antibactériennes contre les agents pathogènes multirésistants (MR) provenant de plaies infectées provenant de trois hôpitaux nigériens.

**Méthodes :** L'extrait de méthanol de la tige de *C. philippensis* a été testé à 20 et 10mg/ml sur 32 pathogènes de plaies comprenant *Staphylococcus aureus* (14), *Pseudomonas aeruginosa* (10) et *Klebsiella pneumoniae* (8). Des méthodes de diffusion d'agar ont été employées pour déterminer l'antibiogramme et la sensibilité aux extraits tandis que la méthode de dilution d'agar a été utilisée pour déterminer les concentrations minimales inhibitrices (CMI) et les concentrations minimales bactéricides (CMB). Le potentiel antioxydant a été évalué par quatre méthodes : activité de catalase, inhibition de peroxydation de lipide, activité de piégeage des radicaux au peroxyde d'hydrogène et l'activité de piégeage des radicaux de DPPH. L'analyse statistique a été effectuée à l'aide d'ANOVA.

**Résultats :** L'antibiogramme a montré que les organismes d'essai étaient des bactéries MR. L'extrait était actif sur tous les agents pathogènes testés avec des zones d'inhibition comprises entre 13-35mm. Les valeurs CMI et CMB varient entre 0,625-5mg/ml pour tous les organismes d'essai et l'indice CMI indique une activité bactéricide. L'extrait de méthanol a montré une bonne activité antioxydante (inhibition de peroxydation lipidique et activité catalase) comparable à la norme d'acide ascorbique.

**Conclusion :** Les activités antioxydantes et antibactériennes de *Celtis philippensis* ont justifié son utilisation folklorique dans le traitement des infections. En outre, une bonne activité sur les pathogènes des plaies multirésistantes met en évidence son potentiel en tant que source d'agents antimicrobiens pour traiter les plaies infectées par des agents pathogènes MR.

**Mots-clés :** *Celtis philippensis*, pathogènes multirésistants aux médicaments, antibactériens, antioxydants, plaies

## INTRODUCTION

Wound infection is one of the most common diseases in developing countries because of poor hygiene causing morbidity and mortality amidst the populace.<sup>1,2</sup> Most community acquired wound infections are polymicrobial in nature of which 25% are caused by *Staphylococcus aureus* while Gram-negative bacteria like *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Peptostreptococcus* are also implicated.<sup>3,4</sup> The risk of developing a wound infection particularly with multi-drug resistant strains of pathogenic bacteria is on the increase.<sup>5</sup> In a 5-year surveillance of wound infections study conducted in Edo State, Nigeria, the overall prevalence of wound infection was 70.1% with *S. aureus* and *P. aeruginosa* as the prevalent organisms all through the years of study.<sup>6</sup> Many studies carried out in different parts of Nigeria showed that *P. aeruginosa*, *S. aureus*, *Klebsiella* species, *E. coli* and *Proteus* species were the most common pathogens isolated from wounds.<sup>7</sup> Today, due to high cost of effective antibiotics and the prevalent antibiotic resistance among microbial strains worldwide, about 60-85% of the population of developing world rely either on herbal or indigenous forms of complementary and alternative medicines for treatment or management of their infections and diseases.<sup>8,9,10</sup> Moreover, free radicals and reactive oxygen species (ROS) have been implicated in many ailments including wound infection. Medicinal plants with antioxidant activity can play a role in the removal or scavenging of toxic free radicals from human body.<sup>11</sup> Medicinal plants have been used since time immemorial for treatment of various ailments of skin and dermatological disorders especially cuts, wounds and burns.<sup>12</sup>

*Celtis philippensis* Blanco (Ulmaceae) also known as *Celtis wightii* Planch in Nigeria, is a small forest tree widely distributed in tropical Africa, Asia and Australia. Among other *Celtis*, it has characteristic leaves that are toothed and prominently 3-nerved;<sup>13</sup> found in Nigeria and commonly distributed across the region from Senegal to Camerouns. Medicinally, its leaf-sap is used in treating cutaneous and subcutaneous parasitic infection and in India to treat diabetes. The genus *Celtis* is of great medicinal importance in the family Ulmaceae (or Cannabaceae); species worthy of note is *Celtis australis* L., (Mediterranean hackberry) whose decoctions of the leaves and fruits were used to astringe the mucous membrane in peptic ulcers, diarrhoea, and dysentery and as a remedy for heavy menstrual bleeding. Also, *Celtis occidentalis* L., of which native Americans used decoctions prepared from the bark as an aid in menses and to relieve sore throat and used the wood extract in

treating jaundice.<sup>14</sup> Others are *Celtis africana* Burm. f, and *Celtis iguana* (Jacq.) Sarg. Due to the immense ethnomedicinal claims and the antimicrobial activities of allied species, we thereby investigated and hereby report the phytochemical constituents, antioxidant and antibacterial activities of *Celtis philippensis* extracts on multidrug resistant wound pathogens from three Nigerian hospitals.

## METHODS

### Plant material

Fresh stems of *Celtis philippensis* were collected from the Botanical Garden of the University of Ibadan. The plant material was identified and authenticated in Forestry Research Institute of Nigeria (FRIN) with a deposited voucher specimen number FHI NO: 109993. The stems were air-dried, milled into coarse powders and extracted with 40% methanol using a Soxhlet extractor. The extract was evaporated under reduced pressure, dried and stored at 4°C for use.

### Test organisms

Thirty-two (32) clinical isolates comprising of *Staphylococcus aureus* (14), *Pseudomonas aeruginosa* (10) and *Klebsiella pneumoniae* (8) from wound infections were collected from Medical Microbiology departments of three Nigerian hospitals; University College Hospital, Ibadan, University of Ilorin Teaching Hospital, Ilorin and Obafemi Awolowo Teaching Hospital, Ile-Ife, Nigeria. The identity of the isolates was confirmed by standard cultural and biochemical tests. Typed strains of *Staphylococcus aureus* (ATCC 25923) and *Pseudomonas aeruginosa* (ATCC 15442) used as reference standards were obtained from Pharmaceutical Microbiology laboratory, University of Ibadan. They were all maintained on agar slants at 4°C prior to use.

### Antibiogram determination

The susceptibility testing of the isolates to standard antibiotics was performed using Kirby-Bauer disc diffusion method. The antibiotics used were selected based on the type of wound pathogen (whether Gram-positive or Gram-negative) and according to drug treatment of infections caused by the pathogen. Therefore, the following antimicrobial agents were used as follows: For *Staphylococcus aureus*, AMX= amoxicillin, OFL= ofloxacin, STR=streptomycin, CEF= ceftriaxone, GEN= gentamicin, PEF= pefloxacin, COT=cotrimoxazole, CPX= ciprofloxacin, ERY= erythromycin, CHL= chloramphenicol; OX= oxacillin, FOX= cefoxitin; for *Pseudomonas aeruginosa*: CXM=cefotaxime,

OFL=ofloxacin, AUG=augmentin, NIT=nitrofurantoin, CPR=ciprofloxacin, CAZ=ceftazidime, CRX=cefuroxime, GEN=gentamicin, IMP= imipenem; for *Klebsiella pneumoniae*: CXM=cefotaxime; OFL=ofloxacin, AUG=augmentin; NIT=nitrofurantoin; CPR=ciprofloxacin, CAZ=ceftazidime; CRX=cefuroxime; GEN=gentamicin; IMP= imipenem; CTR= ceftriazone. The zones of inhibition around each disc was recorded and interpreted by CLSI guidelines.<sup>21</sup>

### Antimicrobial screening

Using agar-well diffusion method as described by Perez *et al.*,<sup>15</sup> a 0.1ml of a two-fold dilution of overnight broth culture of each bacterium (containing  $1.0 \times 10^8$  cfu/ml, equivalent to the turbidity of 0.5 McFarland standard) was seeded into 20ml nutrient agar. A sterile cork borer (diameter 8mm) was used to punch uniform wells on the set and dried agar. Each well was filled with 0.2ml of the crude extracts. Control wells containing gentamicin (10ug/ml), Ceftriaxone (30 ug/ml) and 40% methanol were used as positive and negative controls respectively. A pre-incubation period of about 1hour at 4°C was allowed for the diffusion of extracts and test solutions before incubation. Bacterial plates were incubated at 37°C for 24hours after which the zones of inhibition were measured.

### Determination of minimum inhibitory concentration (MIC) of the plant extracts

The minimum inhibitory concentration of the methanolic plant extracts were determined by the agar dilution method. The plant extracts were prepared in graduated decreasing serial two-fold dilutions to produce the following concentrations: 20, 10, 5, 2.5, 1.25, 0.625, 0.312 and 0.156mg/ml. Each bottle containing 18ml of nutrient agar was melted and cooled to 45-50°C and 2ml of each of the concentration or dilution was added. The bottle was mixed properly and poured in a Petri dish. The agar was then allowed to set firmly. The organisms were inoculated by streaking on the plates which were then incubated at 37°C for 24 hours. The MIC was taken as the minimum concentration of plant extracts that inhibited discernible bacterial growth in the plates.<sup>16</sup>

### Determination of minimum bactericidal concentration (MBC) of the plant extracts

The MBC was determined for all the isolates by sub-culturing from all the test mixtures that failed to show growth in the tube of MIC on Nutrient agar plates using a sterile wire-loop and incubated at 37°C for 24 hours. The MBC was recorded as the lowest concentration of antibiotic inhibiting all growth.<sup>16</sup>

## Determination of antioxidant potentials

### Catalase activity

The activity of catalase (CAT) was measured according to the method of Aebi,<sup>17</sup> as the disappearance of hydrogen peroxide at 240 nm in a reaction medium containing 1800 µl of 50 mM phosphate buffer (pH 7.0). 180 µl of 300 mM H<sub>2</sub>O<sub>2</sub>, and 20 µl of methanolic extract (1:50 dilution). The reaction was monitored for 2 mins (10 intervals), at 240 nm using a UV-visible spectrophotometer (Cecil CE 7200 spectrophotometer, Cecil instrument limited, Milton Technical Centre, England) and expressed as µmol of H<sub>2</sub>O<sub>2</sub> consumed/min/mg extract).

### DPPH radical scavenging activity

The methanolic extract (100 µl, 1mg/ml) was added to 3.9ml of DPPH solution (0.025g/L) and the reactants were incubated at 25°C for 30 mins. Instead of extract, a positive control of AsA was used. The mixture was shaken and allowed to stand in the dark at room temperature for 35 mins. Blank methanol and ascorbic acid (25 - 0.78 µg/ml) were treated in the same way and served as negative and positive controls respectively. Free radical scavenging activity was calculated from absorbance values at 517nm<sup>18</sup> using the UV-spectrophotometer (Cecil CE 7200 spectrophotometer, Cecil instrument limited, Milton Technical Centre, England). The percentage reduction of DPPH was calculated using the following equation: DPPH scavenging activity (%) = [(A0-A1)/A0] x 100. Where A0 = absorbance of negative control, A1 = absorbance of different concentrations of extract or standard drug using the formula: % inhibition (radical scavenging %) = (AC-AS) / Ac x 100.

### Hydrogen peroxide radical scavenging (HPRS) activity

One millimetre of extract (250µg/ml) was mixed with 2.4ml of 0.1 M phosphate buffer (pH 7.4) and then 0.6 of a 43Mm solution of H<sub>2</sub>O<sub>2</sub> in the same buffer were added.<sup>19</sup> After 40 mins, the absorbance of reaction mixture was taken at 230nm against a blank solution (phosphate buffer without H<sub>2</sub>O<sub>2</sub>) using the UV-spectrophotometer (Cecil CE 7200 spectrophotometer, Cecil instrument limited, Milton Technical Centre, England). Percentage scavenging of H<sub>2</sub>O<sub>2</sub> was calculated with Ascorbic acid as control.

$$\% \text{ inhibition (radical scavenging \%)} = (\text{AC-AS}) / \text{Ac} \times 100$$

### Lipid peroxidation inhibition (LPI) using thiobarbituric acid method

Extracts (2ml) and standard solutions (2ml) of the extract

were added to 1ml of 20% aqueous trichloroacetic acid and 2 ml of 0.67% aqueous thiobarbituric acid. After boiling for 10mins, the samples were cooled. The tubes were centrifuged at 3000 rpm for 30 min. Absorbance of the supernatant was measured at 532 nm in a spectrophotometer 20 (Cecil CE 7200 spectrophotometer, Cecil instrument limited, Milton Technical Centre, England). Percentage scavenging of H<sub>2</sub>O<sub>2</sub> was calculated with Ascorbic acid as control.

% inhibition (radical scavenging %) =  $(AC-AS) / AC \times 100$ .

AS= Absorbance of the control; AC= Absorbance of the sample.

### Statistical analysis

Data were analyzed using Analysis of Variance (ANOVA) at 1% level of significance. A difference was considered statistically significant at  $p < 0.05$  (R programme was used).

### RESULTS

The antibiogram of the clinical isolates of the test bacteria showed the resistance levels of the organisms as shown in Tables 1, 2 and 3 for *Staphylococcus aureus*, *Pseudomonas aureus* and *Klebsiella pneumoniae*, respectively. For ease of comparison and interpretation, the percentage resistance to each antimicrobial agent is presented as a summary in Table 4. All the zones of inhibition of the antimicrobial agents on each bacteria strain were classified into Resistance<sup>®</sup>, Intermediate (I)

and Susceptible (S) according to CLSI, 2007.<sup>21</sup> The in vitro antimicrobial screening of the methanolic extracts on *Staphylococcus aureus*, *Pseudomonas aureus* and *Klebsiella pneumoniae* isolates used in this study were recorded as shown in Table 5. Both concentrations used (20mg/ml and 10mg/ml) of the methanolic extracts showed a clear zone of inhibitions comparable to those shown by the positive controls (ceftriaxone and gentamicin). The diameter of zone of inhibition ranged between 16-35mm at 20mg/ml and 13-25mm at 10mg/ml. The minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC) of the methanolic extract (shown in Table 6) ranged between 0.625-5 mg/ml for *K. pneumoniae* isolates, 0.625mg/ml for *P. aeruginosa* isolates and 1.25-2.5mg/ml for *S. aureus* isolates. The minimum bactericidal concentration of the methanolic extracts varies between 1.25-5mg/ml for *S. aureus* and *K. pneumoniae* isolates, 0.625-5mg/ml for *P. aeruginosa* isolates. The extract had a higher MIC values for the *S. aureus* test isolates than that for *K. pneumoniae* and *P. aeruginosa* test isolates. The result of the Analysis of variance (ANOVA) shows that there is a significance difference among the variables measured at 1% level of significance on the *P. aeruginosa* isolates but no significance difference among the *S. aureus* and *K. pneumoniae* isolates when compared by Tukey's post-test in a One-way ANOVA analysis. The results of three antioxidant assays were presented in Figure 1 as a bar chart in comparison with ascorbic acid control.

## Antibiogram of clinical isolates from wound infection

Table 1: Antibiotic susceptibility of *Staphylococcus aureus* Isolates

Isolate/Abs	AMX	OFL	STR	CHL	CEF	GEN	PEF	COT	CPX	ERY	OX	FOX
S1	R	R	I	S	I	R	I	R	I	R	R	R
S2	R	R	R	R	I	R	I	R	R	R	R	R
S3	R	S	I	I	R	R	S	R	S	R	R	R
S4	R	R	R	I	I	R	I	R	I	R	R	R
S5	R	R	I	R	R	I	R	R	R	S	R	R
S6	R	R	R	R	I	R	R	I	R	R	R	R
S7	R	R	R	R	R	R	R	R	R	R	R	R
S8	R	R	R	R	I	R	R	R	R	R	R	R
S9	R	R	R	R	I	R	R	R	R	R	R	R
S10	R	I	S	I	I	R	R	R	I	R	R	R
S11	R	R	R	R	S	R	R	R	R	R	R	S
S12	R	S	S	R	R	R	I	R	I	R	R	R
S13	R	S	R	R	R	R	R	R	I	R	R	R
S14	R	S	S	S	S	S	S	S	I	R	R	R

Key: R= Resistant; AMX= amoxicillin, OFL= ofloxacin, STR=streptomycin, CEF= ceftriaxone, GEN= gentamicin, PEF= pefloxacin, COT=cotrimoxazole, CPX= ciprofloxacin, ERY= erythromycin, CHL= chloramphenicol; OX= oxacillin, FOX= cefoxitin

**Table 2: Antibiotic susceptibility of *Pseudomonas aeruginosa* isolates**

Isolate	CXM	OFL	AUG	NIT	CPR	CAZ	CRX	GEN	IMP
P1	R	S	R	R	S	S	R	S	R
P2	R	R	R	R	R	R	R	R	R
P3	R	R	R	R	R	R	R	R	R
P4	R	R	R	R	R	S	R	R	R
P5	R	R	R	R	R	S	R	R	R
P6	R	R	R	R	R	R	R	R	R
P7	R	R	R	R	R	R	R	R	R
P8	R	R	R	R	R	R	R	R	R
P9	R	R	R	R	R	R	R	R	R
P10	R	S	R	R	S	S	R	S	R

Key: R=Resistant, CXM=cefotaxime, OFL=ofloxacin, AUG=augmentin, NIT=nitrofurantoin, CPR=ciprofloxacin, CAZ=ceftazidime, CRX=cefuroxime, GEN=gentamicin, IMP= imipenem

**Table 3: Antimicrobial susceptibility of *Klebsiella pneumoniae* isolates**

ISOLATES	CRX	CPR	NIT	AUG	OFL	CXM	GEN	CTR	IMP	CAZ
K2	R	R	R	R	I	R	R	R	S	R
K3	R	R	R	R	S	R	R	I	S	I
K4	R	I	S	S	S	I	S	I	S	S
K5	R	S	R	R	I	R	S	R	I	R
K6	R	I	I	R	I	R	S	R	R	R
K8	R	R	S	R	R	R	R	I	S	R
K16	R	S	R	R	S	R	S	R	S	S
K18	R	R	R	R	R	R	S	R	R	R

Key: R=Resistant; I=Intermediate; S=Sensitive; CXM=cefotaxime; OFL=ofloxacin, AUG=augmentin; NIT=nitrofurantoin; CPR=ciprofloxacin, CAZ=ceftazidime; CRX=cefuroxime; GEN=gentamicin; IMP= imipenem; CTR=ceftriazone

**Table 4: Summary of antimicrobial resistance of *Staphylococcus*, *Pseudomonas* and *Klebsiella* species**

	<i>Pseudomonas aeruginosa</i>	<i>Klebsiella pneumonia</i>	<i>Staphylococcus aureus</i>
Antibiotics	Resistance (%)	Resistance (%)	Resistance (%)
Cefotaxime	100.00	87.50	—
Augmentin	100.00	87.50	—
Nitrofurantoin	100.00	62.50	—
Cefuroxime	100.00	100.00	—
Ceftazidime	60.00	62.50	—
Imipenem	100.00	25.00	—
Gentamicin	80.00	37.50	85.71
Ofloxacin	80.00	25.00	64.29
Ciprofloxacin	80.00	—	50.00
Ceftriaxone	—	62.50	35.71
Pefloxacin	—	—	57.14
Streptomycin	—	—	57.14
Erythromycin	—	—	92.86
Cotrimoxazole	—	—	85.71
Chloramphenicol	—	—	64.29
Amoxicillin	—	—	100.00
Oxacillin	—	—	100.00
Cefoxitin	—	—	92.86

**Key: — = Not tested**



**Table 5.0: Antimicrobial activity of *Celtis philippensis* stem extract on wound pathogens**

Isolates	Diameter(mm) of Zone of inhibition of Extract/Controls on Test Bacteria			
	<i>Celtis philippensis</i> stem extract		Gentamicin	Ceftriaxone
	20mg/ml	10mg/ml	10 µg	30 µg
S1	19	14	23	25
S2	17	14	25	1,18
S3	20	16	24	30
S4	20	16	25	30
S5	21	15	15	1,20
S6	20	13	R,8	1,18
S7	20	18	R,8	20
S8	18	17	R,8	1,15
S9	20	20	R8	R,8
S10	25	25	20	30
S11	24	22	32	1,20
S12	27	25	26	32
S13	18	16	30	25
S14	30	27	25	1,20
P1	22	20	23	1,16
P2	21	18	R,8	R,8
P3	22	20	R,8	R,8
P4	25	20	R,8	R,8
P5	25	20	24	1,18
P6	28	25	R,8	R,8
P7	24	22	R,8	R,8
P8	22	19	R,8	R,8
P9	26	20	R,8	1,20
P10	26	22	17	25
K2	20	16	20	30
K3	20	16	20	22
K4	26	16	20	24
K5	28	20	21	26
K6	20	16	1,14	1,20
K8	16	14	22	20
K16	35	22	20	30

**Keys:** S1 – S14 = *Staphylococcus aureus*, P1 – P10 = *Pseudomonas aeruginosa*, K2 – K16 = *Klebsiella pneumoniae*

**Table 6.0: MIC and MBC of *Celtis philippensis* extract on test 4bacteria**

ISOLATES	MIC (mg/ml)	MBC (mg/ml)
S1	1.25	2.50
S2	2.50	5.00
S3	2.50	5.00
S6	2.50	5.00
S7	1.25	1.25
S8	2.50	5.00
S9	2.50	5.00
S10	2.50	5.00
P1	0.63	0.63
P2	0.63	2.50
P3	0.63	5.00
P6	0.63	5.00
P7	0.63	5.00
P8	0.63	2.50
P9	0.63	2.50
P10	0.63	2.50
K3	0.63	2.50
K4	0.63	2.50
K5	0.63	5.00
K6	1.25	5.00
K8	5.00	2.50
K16	0.63	1.25
K18	2.50	2.50

**Keys:** S1 – S10 = *Staphylococcus aureus*, P1 – P10 = *Pseudomonas aeruginosa*, K3 – K18 = *Klebsiella pneumoniae*




Fig. 1: Antioxidant activities (%) of *Celtis philippensis* by HPRS, LPI and DPPH methods compared with that of ascorbic acid reference standard

## DISCUSSION

The antibiogram showed that most of the tested wound pathogens were multidrug resistant (MDR), having shown resistance to three or more antimicrobial agents belonging to different classes. The *Pseudomonas* isolates showed 100% resistance to augmentin, cefuroxime, ceftriaxone, nitrofurantoin and imipenem; the most active drug being ceftazidime (40%) and fluoroquinolones (20%). Actually, 60% of the *Pseudomonas* showed 100% resistance to all tested antibacterials. These type of MDR *Pseudomonas aeruginosa* strains were also reported for Egyptian hospitals by Ahmed *et al.*,<sup>22</sup> but this is at higher levels, which is an indication of increasing resistance. In this study, the most active drug on the *Pseudomonas* isolates was ceftazidime which is similar to that reported by Diab *et al.*,<sup>23</sup> here the activity of ceftazidime was 40% while theirs was 54%,<sup>23</sup> which also indicated increasing resistance. The *Klebsiella* strains were relatively more susceptible to the chemotherapeutic agents, but with tendency for more to acquire resistance with time. The *Staphylococci* also showed a high-level resistance which is consistent with that obtainable with methicillin resistant *Staphylococcus aureus* (MRSA), vancomycin intermediate *Staphylococcus aureus* (VISA) and vancomycin resistant *Staphylococcus aureus* (VRSA). The *Staphylococcus aureus* strains were 93-100% resistant to the  $\beta$ -lactam drugs but were most susceptible to ceftriaxone 35%. As MRSA is the commonest aetiology of nosocomial wound infection,<sup>[24]</sup> its eradication has become more difficult with the emergence of VISA and VRSA. The antibiogram has clearly revealed that the wound pathogens used were actually MDR which enables us to appreciate the significant broad-spectrum activity of the plant's extracts on the pathogens.<sup>24</sup>

The role of medicinal herbs in treating wound infection dated from Biblical era when extract from Fig tree, *Ficus carica* (Moraceae) was used to heal the boil on king Hezekiah.<sup>25</sup> With increasing antimicrobial resistance worldwide, the search for alternative therapy among natural herbs is essential and must be taken from empirical to scientific documentation. In this study, it behoves us to report that the extract of *Celtis philippensis* was very active on MDR wound pathogens. The Gram-negative bacteria were more susceptible to the plant extracts than Gram-positive bacteria which contradicts some previous reports that plant extracts were more active against Gram-positive bacteria than Gram-negative bacteria.<sup>26,27</sup> *Pseudomonas aeruginosa* isolates which were highly resistant to the antimicrobials were highly susceptible to the extract. The MDR

*Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Klebsiella pneumonia* were all susceptible to the extract at 10 and 20 mg/ml. The minimum inhibitory concentration (MIC) of the extract that inhibited the antibiotic resistant organisms are 0.625mg/ml (for *Pseudomonas* and four of the *Klebsiella*) and 2.5mg/ml for the *S. aureus* which were the least susceptible. MIC Index of the extract (MBC/MIC) was less than 4 indicating bactericidal activity especially of the stem extract.<sup>28</sup> The *S. aureus* test isolates were not very susceptible to the extract. Ability of herbal extracts to overcome resistant pathogens can be attributed to the presence of multiple antimicrobial compounds, synergistic or additive effect among bioactive compounds and diverse nature of constituents that organisms cannot readily produce genes against.

The roles of antioxidants in wound healing are well documented. Natural antioxidants in medicinal plants occur in form of phenolics, flavonoids and polyhydroxyl compounds which are able mop-up or scavenge the reactive radicals and oxygen species, thereby enhancing tissue repair and promote wound healing. According to Dudonnei *et al.*,<sup>29</sup> antioxidant activities of medicinal plants must be evaluated by at least two methods in order to take into account different modes of action of a given antioxidant.<sup>29</sup> The antioxidant activities revealed that the extract showed high potential in the lipid peroxidation inhibition and catalase activity. Its activity on hydrogen peroxide scavenging and DPPH radical scavenging were relatively low compared with the ascorbic acid standard. Phytochemicals present in the extract has shown the ability to scavenge hydroxyl radical that reacts with the methylene groups of polyunsaturated fatty acids (PUFA), the main components of membrane lipids, are susceptible to peroxidation and thus preventing the aldehydes (final stable products of per-oxidation) to react with TBA to form thiobarbituric acid-malonaldehyde adduct with an absorbance maximum at 532 nm. This led to reduction of adduct formation indicating their lipo-protective potential. The extract also showed the capability to catalyze the dismutation of hydrogen peroxide in water and oxygen. The antimicrobial activity study could not determine the mechanism involved in the bactericidal potency of the extracts on MDR wound pathogens used in this work.

## CONCLUSION

The extract was active against both Gram-positive and Gram-negative bacterial isolates tested and this may indicate a broad spectrum of activity. The results of the

study support the traditional application of *C. Philippensis* in treating bacterial, cutaneous and subcutaneous parasitic infections. The susceptibility of all the test organisms (MDR), including *P. aeruginosa* that has resistance against many antibiotics goes further to prove that the plant has potentials as alternative source of antimicrobial agent in overcoming the problems of resistance to chemotherapeutic drugs. *Celtis philippensis* stem has also been revealed as a source of antioxidant agents. It is therefore desirable to isolate and characterize the antimicrobial and antioxidant agents from *C. philippensis*.

#### ACKNOWLEDGEMENT

The authors appreciate the roles of Mr Kayode Owolabi of Botany Department, University of Ibadan in identifying the plant, Mr Samuel Odewor of FRIN for authentication; and the Staff of Biochemistry and Pharmaceutical Microbiology laboratories of University of Ibadan for technical assistance.

#### REFERENCES

1. Kumar MS, Sripriya R, Raghavan HV, Sehgal PK (2006). Wound healing potential of *Cassia fistula* on infected albino rat model. *Journal of Surgical Research* 131: 283-289.
2. Akingbade OA, Akinjinmi AA, Ezechukwu US, Okerentugba PO, Nwanze JC, Onoh CC, Okonko IO (2013). Antibacterial effect of *Moringa oleifera* on multidrug resistant *Pseudomonas aeruginosa* isolates from wound infections in Abeokuta, Ogun State, Nigeria. *World Rural Observation* 5(3):6-10.
3. Bowler P, Duerden B and Armstrong D (2001). Wound microbiology and associated approaches to wound management. *Clinical Microbiology Review* 14(2): 244-69.
4. Posluszny J, Conrad P and Marcia H (2011). Surgical burn wound infections and their clinical implications. *Journal of Burn Care Research* 32(2): 324-33.
5. Motayo BO, Akinbo JA, Ogiogwa IJ, Idowu AA, Nwanze JC, Onoh CC and Okerentugba PO (2013). Innocent-Adiele H. C, Okonko, I. O. Bacteria Colonisation and Antibiotic Susceptibility Pattern of Wound Infections in a Hospital in Abeokuta. *Frontiers in Science* 3 (1): 43-48.
6. Oladeinde BH, Omoregie R, Olley M, Anunibe JA and Onifade AA (2012). A 5 - year surveillance of wound infections at a rural tertiary hospital in Nigeria. *African Health Science* 13 (2): 351-356.
7. Akingbade OA, Balogun SA, Ojo DA, Afolabi RO, Motayo BO, Okerentugba PO and Okonko IO (2012). Plasmid Profile Analysis of Multidrug Resistant *Pseudomonas aeruginosa* Isolated from Wound Infections in South West, Nigeria. *World Applied Sciences Journal* 20(6): 766-775.
8. Okeke IN, Lamikanra A and Edelman R (1996). Socioeconomic and behavioural factors leading to acquired bacterial resistance to antibiotics in developing countries. *Emerging Infectious Diseases* 5: 18-27.
9. Archana SJ, Paul R and Tiwari A (2011). Indian medicinal plants: A rich source of natural immunomodulator. *International Journal of Pharmacology* 7(2): 198-205.
10. Mahima RA, Deb R, Latheef S and Samad HA (2012). Immunomodulatory and therapeutic potentials of herbal, traditional/indigenous and ethnoveterinary medicines. *Pak.J.Biol.Sci.* 15: 754-774.
11. Dong-Ping X, Ya L, Xiao M, Tong Z, Yue Z, Jie Z, Jiao-Jiao Z and Hua-Bin L (2017). Natural antioxidants in foods and medicinal plants: extraction, assessment and resources. *International Journal of Molecular Science* 18(1): 96
12. Kumar B, Vijayakumar M, Govindarajan R and Pushpangadan P (2007). Ethnopharmacological approaches to wound healing-exploring medicinal plants of India. *Journal of Ethnopharmacology* 114: 103-113
13. Keay, R.W.J. (1989). *Trees of Nigeria*. Oxford University Press. Pg 282-287
14. El-Alfy TS, El-Gohary HMA, Sokkar NM, Horsny M and Al-Mahdy DA (2011). A new flavonoid C-Glycoside from *Celtis australis* L. and *Celtis occidentalis* L. leaves and potential antioxidant and cytotoxic activities. *Scientia Pharmaceutica* 79(4): 963-975.
15. Perez C, Pauli M and Bazerque P (1990). An antibiotic assay by agar-well diffusion method. *Acta Biologica et Medicinæ Experimentalis* 15: 113-115
16. Cheesebrough, M (2000). *District laboratory practice in tropical countries Part 2*. UK: Cambridge University Press. 134-143
17. Aebi H (1984). Catalase. In: *Methods of Enzymatic Analysis*, Bergmeyer, H.U. (Ed.). Academic Press Inc., New York, USA. 2: 673-685.
18. Wettasinghe M and Shahidi F (2000). Scavenging of reactive oxygen species and DPPH free radicals by extracts of borage and evening primrose meals. *Food Chemistry* 70 (1): 17-26.

19. Sowndharrajan K, Siddhuraju P and Manian S (2010). *In vitro* evaluation of the antioxidant activities in the differentially processed seeds from underutilized legume, *Bauhinia vahlii* Wight & Arn. *Food Science and Biotechnology* 19: 503-509.
20. Saha K, Lajis NH, Israif DA, Hamzah AS, Khozirah S, Khamis S, Syahida A (2004). Evaluation of antioxidant and nitric oxide inhibitory activities of selected Malaysian medicinal plants. *Journal of Ethnopharmacology* 92(2-3): 263-267.
21. Clinical Laboratory Standard Institute [CLSI] (2007). Performance Standard for Antimicrobial Susceptibility Testing. Seventeenth Informational Supplement. M100-S17. Wayne, Pennsylvania, USA.
22. Ahmed FY, Farghaly Aly U, Abd El-Baky RM and Waly NGFM (2020). Comparative Study of Antibacterial Effects of Titanium Dioxide Nanoparticles Alone and in Combination with Antibiotics on MDR *Pseudomonas aeruginosa* Strains. *International Journal of Nanomedicine* 15: 3393-3404.
23. Diab M, Fam N, El-said M, El-dabaa E, El Defrawy I and Saber M (2013). Occurrence of VIM-2 metallo- $\beta$ -lactamases in imipenem resistant and susceptible *Pseudomonas aeruginosa* clinical isolates from Egypt. *African Journal of Microbiology Research* 7(35): 4465-4472.
24. Dou JL, Jiang YW, Xie JQ and Zhang XG (2016). New Is Old, and Old Is New: Recent Advances in Antibiotic-Based, Antibiotic-Free and Ethnomedical Treatments against Methicillin-Resistant *Staphylococcus aureus* Wound Infections. *International Journal of Molecular Science* 17(5): 617.
25. King James Bible (2000). The Zondervan Corporation. 2Kings 20(7): pp 388
26. Rabe T and Van Staden J (1997). Antibacterial activity of South African plants used for medicinal purposes. *Journal of Ethnopharmacology* 56: 81-87.
27. Vlietinck AJ, Van Hoof L, Totte J, Lasure A, Vanden Berghe D, Rwangobo PC and Mvukiyuniwami J (1995). Screening of hundred Rwandese medicinal plants for antimicrobial and antiviral properties. *Journal of Ethnopharmacology* 46: 31-47.
28. Mamidou Koné W, Kamanzi Atindehou K, Kacou-N'Douba A and Dosso M (2007). Evaluation of 17 medicinal plants from northern Côte d'Ivoire for their *in vitro* activity against *Streptococcus pneumoniae*. *African Journal of Traditional, Complementary and Alternative Medicines* 4(1): 17-22.
29. Dudonné S, Vitrac X, Coutière P, Woillez M and Meïrillon JM (2009). Comparative study of antioxidant properties and total phenolic content of 30 plant extracts of industrial interest using DPPH, ABTS, FRAP, SOD, and ORAC assays. *Journal of Agricultural and Food Chemistry* 57:1768-1774